



Carrum Primary School 3385

2-6 Church Road Carrum VIC 3197

PO Box 156 Seaford VIC 3198Ph:

(03) 9772 1117

carrum.ps@education.vic.gov.au

ANAPHYLAXIS POLICY 2026



Help for non-English speakers

If you need help to understand the information in this policy, please contact school administration on 03 9772 1117.

PURPOSE

To explain to Carrum Primary School parents, carers, staff and students the processes and procedures in place to support students diagnosed as being at risk of suffering from anaphylaxis. This policy also ensures that Carrum Primary School is compliant with Ministerial Order 706 and the department's guidelines for anaphylaxis management.

SCOPE

This policy applies to:

- all staff, including casual relief staff, canteen operators and volunteers
- all students who have been diagnosed with anaphylaxis, or who may require emergency treatment for an anaphylactic reaction, and their parents and carers.

POLICY

School Statement

Carrum Primary School will fully comply with Ministerial Order 706 and the associated guidelines published by the Department of Education.

Anaphylaxis

Anaphylaxis is a severe allergic reaction that occurs after exposure to an allergen. The most common allergens for school-aged children are nuts, eggs, cow's milk, fish, shellfish, wheat, soy, sesame, latex, certain insect stings and medication.

Symptoms

Signs and symptoms of a mild to moderate allergic reaction can include:

- swelling of the lips, face and eyes
- hives or welts
- tingling in the mouth.

Signs and symptoms of anaphylaxis, a severe allergic reaction, can include:

- difficult/noisy breathing

- swelling of tongue
- difficulty talking and/or hoarse voice
- wheeze or persistent cough
- persistent dizziness or collapse
- student appears pale or floppy
- abdominal pain and/or vomiting.

Symptoms usually develop within ten minutes and up to two hours after exposure to an allergen but can appear within a few minutes.

Treatment

Adrenaline given as an injection into the muscle of the outer mid-thigh is the first aid treatment for anaphylaxis.

Individuals diagnosed as being at risk of anaphylaxis are prescribed an adrenaline device for use in an emergency. These adrenaline devices are designed so that anyone can use them in an emergency.

Individual Anaphylaxis Management Plans

All students who are diagnosed by a medical practitioner as being at risk of suffering from an anaphylactic reaction must have an Individual Anaphylaxis Management Plan. When notified of an anaphylaxis diagnosis, the Principal (or delegate) of Carrum Primary School is responsible for developing a plan in consultation with the student's parents/carers.

Where necessary, an Individual Anaphylaxis Management Plan will be in place as soon as practicable after a student enrolls at Carrum Primary School and where possible, before the student's first day.

Parents and carers must:

- obtain an ASCIA Action Plan for Anaphylaxis (RED) from the student's medical practitioner and provide a copy to the school as soon as practicable
- immediately inform the school in writing if there is a relevant change in the student's medical condition and obtain an updated ASCIA Action Plan for Anaphylaxis (RED)
- provide an up-to-date photo of the student for the ASCIA Action Plan for Anaphylaxis (RED) and provide it to the school each time it is updated
- provide the school with a current adrenaline device for the student that has not expired
- participate in annual reviews of the student's Individual Anaphylaxis Management Plan that is prepared by the school.

Each student's Individual Anaphylaxis Management Plan must include:

- information about the student's allergies and the potential for anaphylactic reaction, including the type of allergies the student has
- information about the signs or symptoms the student might exhibit in the event of an allergic reaction based on a written diagnosis from a medical practitioner
- strategies to minimise the risk of exposure to known allergens while the student is under the care or supervision of school staff, including in the school yard, at camps and excursions, or at special events conducted, organised or attended by the school
- the name of the person(s) responsible for implementing the risk minimisation strategies, which have been identified in the Individual Anaphylaxis Management Plan
- information about where the student's medication will be stored
- the student's emergency contact details
- an up-to-date ASCIA Action Plan for Anaphylaxis (RED) completed by the student's medical practitioner.

Review and updates to Individual Anaphylaxis Management Plans:

A student's Individual Anaphylaxis Management Plan will be reviewed and updated on an annual basis in consultation with the student's parents/carers. The plan will also be reviewed and, where necessary, updated in the following circumstances:

- as soon as practicable after the student has an anaphylactic reaction at school
- if the student's medical condition, insofar as it relates to allergy and the potential for anaphylactic reaction, changes
- when the student is participating in an off-site activity, including camps and excursions, or at special events including fetes and concerts.

Our school may also consider updating a student's Individual Anaphylaxis Management Plan if there is an identified and significant increase in the student's potential risk of exposure to allergens at school.

Location of plans and adrenaline devices

A copy of each student's Individual Anaphylaxis Management Plan will be stored with their ASCIA Action Plan for Anaphylaxis (RED) in the student's classroom, together with the student's adrenaline device. Adrenaline devices must be labelled with the student's name. Copies of each student's Individual ASCIA plans are also kept in the First Aid room with a school general use Adrenaline autoinjector available where required.

Risk Minimisation Strategies

To reduce the risk of a student suffering from an anaphylactic reaction at Carrum Primary School, we have put in place the following strategies:

	CLASSROOM
1.	Keep a copy of the student's Individual Anaphylaxis Management Plan in the classroom. Be sure the ASCIA Action Plan for Anaphylaxis is kept with the student's adrenaline autoinjector.
2.	Liaise with parents about food-related activities well ahead of time.
3.	Use non-food treats where possible, but if food treats are used in class, it is recommended that parents of students with food allergy provide a treat box with alternative treats. Alternative treat boxes should be clearly labelled and only handled by the student.
4.	Never give food from outside sources to a student who is at risk of anaphylaxis.
5.	Products labelled 'may contain traces of nuts' should not be served to students allergic to nuts. Products labelled 'may contain milk or egg' should not be served to students with milk or egg allergy and so forth.
6.	Be aware of the possibility of hidden allergens in food and other substances used in cooking, food technology, science and art classes (e.g., egg or milk cartons, empty peanut butter jars).
7.	Ensure all cooking utensils, preparation dishes, plates, and knives and forks etc are washed and cleaned thoroughly after preparation of food and cooking.
8.	Children with food allergy need special care when doing food technology. The Kitchen specialist will meet with parents to discuss plan and suitable cooking methods for that student prior to taking on these classes.
9.	Have regular discussions with students about the importance of washing hands, eating their own food and not sharing food.
10.	A designated staff member should inform casual relief teachers, specialist teachers and volunteers of the names of any students at risk of anaphylaxis, the location of each student's Individual Anaphylaxis Management Plan and adrenaline autoinjector, the school's Anaphylaxis Management Policy, and each individual person's responsibility in managing an incident. i.e. seeking a trained staff member.
	YARD

1.	If a school has a student who is at risk of anaphylaxis, sufficient school staff on yard duty must be trained in the administration of the adrenaline autoinjector (i.e. EpiPen®) and be able to respond quickly to an allergic reaction if needed.
2.	The adrenaline autoinjector and each student's individual ASCIA Action Plan for Anaphylaxis must be easily accessible from the yard, and staff should be aware of their exact location. (Remember that an anaphylactic reaction can occur in as little as a few minutes). Where appropriate, an adrenaline autoinjector may be carried in the school's yard duty bag.
3.	Schools must have an emergency response procedure in place so the student's medical information and medication can be retrieved quickly if a reaction occurs in the yard. This may include all yard duty staff carrying emergency cards in yard-duty bags, walkie talkies or yard-duty mobile phones. All staff on yard duty must be aware of the school's emergency response procedures and how to notify the general office/first aid team of an anaphylactic reaction in the yard.
4.	Yard duty staff must also be able to identify, by face, those students at risk of anaphylaxis. Tags with photographs are attached to each yard duty bag to identify students with Anaphylaxis.
5.	Students with severe allergies to insects should be encouraged to stay away from water or flowering plants. School staff should liaise with parents to encourage students to wear light or dark rather than bright colours, as well as closed shoes and long-sleeved garments when outdoors.
6.	Keep lawns and clover mowed and outdoor bins covered.
7.	Students should keep drinks and food covered while outdoors.

	SPECIAL EVENTS (sporting events, incursions, class parties etc)
1.	If a school has a student at risk of anaphylaxis, sufficient school staff supervising the special event must be trained in the administration of an adrenaline autoinjector to be able to respond quickly to an anaphylactic reaction if required.
2.	School staff should avoid using food in activities or games (where practical), including as rewards.
3.	For special events involving food, school staff should consult parents in advance to either develop an alternative food menu or request the parents to send a meal for the student.
4.	Parents of other students should be informed in advance about foods that may cause allergic reactions in students at risk of anaphylaxis and request that they try to avoid providing students with treats whilst they are at school or at a special school event.
5.	Party balloons should not be used if any student is allergic to latex.
6.	If students from other schools are participating in an event at your school, consider requesting information from the participating schools about any students who will be attending the event who are at risk of anaphylaxis. Agree on strategies to minimise the risk of a reaction while the student is visiting the school. This should include a discussion of the specific roles and responsibilities of the host and visiting school. Students at risk of anaphylaxis should bring their own adrenaline autoinjector with them to events outside their own school.

	OUT OF SCHOOL SETTINGS (excursions, sporting events)
1.	If a school has a student at risk of anaphylaxis, sufficient school staff supervising the special event must be trained in the administration of an adrenaline autoinjector and be able to respond quickly to an anaphylactic reaction if required.
2.	A school staff member or team of school staff trained in the recognition of anaphylaxis and the administration of the adrenaline autoinjector must accompany any student at risk of anaphylaxis on field trips or excursions.
3.	School staff should avoid using food in activities or games, including as rewards.
4.	The adrenaline autoinjector and a copy of the individual ASCIA Action Plan for Anaphylaxis for each student at risk of anaphylaxis should be easily accessible and school staff must be aware of their exact location.
5.	For each excursion etc, a risk assessment should be undertaken for each individual student attending who is at risk of anaphylaxis. The risks may vary according to the

	number of anaphylactic students attending, the nature of the excursion/sporting event, size of venue, distance from medical assistance, the structure of excursion and corresponding staff-student ratio. All school staff members present during the excursion need to be aware of the identity of any students attending who are at risk of anaphylaxis and be able to identify them by face.
6.	The school should consult parents of anaphylactic students in advance to discuss issues that may arise, for example to develop an alternative food menu or request the parents provide a special meal (if required).
7.	Parents may wish to accompany their child on field trips and/or excursions. This should be discussed with parents as another strategy for supporting the student who is at risk of anaphylaxis.
8.	Prior to the excursion taking place school staff should consult with the student's parents and medical practitioner (if necessary) to review the student's Individual Anaphylaxis Management Plan to ensure that it is up to date and relevant to the particular excursion activity.
9.	If the field trip, excursion or special event is being held at another school then that school should be notified ahead of time that a student at risk of anaphylaxis will be attending, and appropriate risk minimisation strategies discussed ahead of time so that the roles and responsibilities of the host and visiting school are clear. Students at risk of anaphylaxis should take their own adrenaline autoinjector with them to events being held at other schools.

	CAMP/REMOTE AREAS
1.	Prior to engaging a camp owner/operator's services the school should make enquiries as to whether the operator can provide food that is safe for anaphylactic students. If a camp owner/operator cannot provide this confirmation in writing to the school, then the school should strongly consider using an alternative service provider. This is a reasonable step for a school to take in discharging its duty of care to students at risk of anaphylaxis.
2.	The camp cook should be able to demonstrate satisfactory training in food allergen management and its implications for food-handling practices, including knowledge of the major food allergens triggering anaphylaxis, cross-contamination issues specific to food allergy, label reading, etc.
3.	Schools must not sign any written disclaimer or statement from a camp owner/operator that indicates that the owner/operator is unable to provide food which is safe for students at risk of anaphylaxis. Schools have a duty of care to protect students in their care from reasonably foreseeable injury, and this duty cannot be delegated to any third party.
4.	Schools should conduct a risk assessment and develop a risk management strategy for students at risk of anaphylaxis while they are on camp. This should be developed in consultation with parents of students at risk of anaphylaxis and camp owners/operators prior to the camp's commencement.
5.	School staff should consult with parents of students at risk of anaphylaxis and the camp owner/operator to ensure that appropriate procedures are in place to manage an anaphylactic reaction should it occur. If these procedures are deemed to be inadequate, further discussions, planning and implementation will need to be undertaken in order for the school to adequately discharge its non-delegable duty of care.
6.	If the school has concerns about whether the food provided on a camp will be safe for students at risk of anaphylaxis, it should raise these concerns in writing with the camp owner/operator and also consider alternative means for providing food for those students.
7.	Use of substances containing known allergens should be avoided altogether where possible.
8.	Camps should be strongly discouraged from stocking peanut or tree nut products, including nut spreads. Products that 'may contain' traces of nuts may be served, but not to students who are known to be allergic to nuts. If eggs are to be used there must be suitable alternatives provided for any student known to be allergic to eggs.
9.	Prior to the camp taking place school staff should consult with the student's parents to review the students Individual Anaphylaxis Management Plan to ensure that it is up to date and relevant to the circumstances of the particular camp.
10	The student's adrenaline autoinjector, Individual Anaphylaxis Management Plan, including the ASCIA Action Plan for Anaphylaxis and a mobile phone must be taken on camp. If

	mobile phone access is not available, an alternative method of communication in an emergency must be considered, e.g. a satellite phone. All staff attending camp should familiarise themselves with the students' Individual Anaphylaxis Management Plans AND plan emergency response procedures for anaphylaxis prior to camp and be clear about their roles and responsibilities in the event of an anaphylactic reaction.
11.	Contact local emergency services and hospitals well before the camp to provide details of any medical conditions of students, location of camp and location of any off-camp activities. Ensure contact details of emergency services are distributed to all school staff as part of the emergency response procedures developed for the camp.
12.	It is strongly recommended that schools take an adrenaline autoinjector for general use on a school camp (even if there is no student who is identified as being at risk of anaphylaxis) as a back-up device in the event of an emergency.
13.	Schools should consider purchasing an adrenaline autoinjector for general use to be kept in the first aid kit and include this as part of the emergency response procedures.
14.	Each student's adrenaline autoinjector should remain close to the student and school staff must be aware of its location at all times.
15.	The adrenaline autoinjector should be carried in the school first aid kit; however, schools can consider allowing students, particularly adolescents, to carry their adrenaline autoinjector on camp. Remember that all school staff members still have a duty of care towards the student even if they do carry their own adrenaline autoinjector.
16.	Students with allergies to insects should always wear closed shoes and long-sleeved garments when outdoors and should be encouraged to stay away from water or flowering plants.
17.	Cooking and art and craft games should not involve the use of known allergens.
18.	Consider the potential exposure to allergens when consuming food on buses and in cabins.

Adrenaline autoinjectors for general use

Carrum Primary School will maintain a supply of two EpiPens (adult) and one EpiPen Junior adrenaline autoinjectors for general use, as a back-up to those provided by parents and carers for specific students, and also for students who may suffer from a first-time reaction at school.

There are currently 4 adrenaline devices approved by the Therapeutic Goods Administration for use in Australia: the EpiPen®, the Anapen®, Jext® and Neffy®. All devices can be used when provided by families for students, however, the principal or allocated staff member can only use EpiPen®, Anapen® or Jext® adrenaline autoinjector for general use. For more information about which autoinjector to purchase for general use, refer to [Adrenaline autoinjectors for general use](#).

Adrenaline autoinjectors for general use will be stored at the First Aid room and labelled "general use".

The principal's delegate/ First Aid Officer is responsible for arranging the purchase of adrenaline autoinjectors for general use, and will consider:

- the number of students enrolled at Example School at risk of anaphylaxis
- the accessibility of adrenaline autoinjectors supplied by parents
- the availability of a sufficient supply of autoinjectors for general use in different locations at the school, as well as at camps, excursions and events
- the limited life span of adrenaline autoinjectors, and the need for general use adrenaline autoinjectors to be replaced when used or prior to expiry
- the weight of the students at risk of anaphylaxis to determine the correct dosage of adrenaline autoinjector/s to purchase.

Emergency Response

In the event of an anaphylactic reaction, the emergency response procedures in this policy must be followed, together with the school's general first aid procedures, emergency response procedures and the student's Individual Anaphylaxis Management Plan.

A complete and up-to-date list of students identified as being at risk of anaphylaxis is maintained by Rebecca Schefman (School Administration) and Cathy Vittas (First Aider) and stored in the First Aid room and in the Administration office. For camps, excursions and special events, a designated staff member will be responsible for maintaining a list of students at risk of anaphylaxis attending the special event, together with their Individual Anaphylaxis Management Plans and adrenaline devices, where appropriate.

If a student experiences an anaphylactic reaction at school or during a school activity, school staff must:

Step	Action
1.	<ul style="list-style-type: none"> • Lay the person flat • Do not allow them to stand or walk • If breathing is difficult, allow them to sit with legs outstretched • Be calm and reassuring • Do not leave them alone • Seek assistance from another staff member or reliable student to locate the student's adrenaline device or the school's general use autoinjector, and the student's Individual Anaphylaxis Management Plan, stored in the student's individual classroom and in the First Aid room. • If the student's plan is not immediately available, or they appear to be experiencing a first-time reaction, follow steps 2 to 5
2.	<p>Administer an EpiPen or EpiPen Jr</p> <ul style="list-style-type: none"> • Remove from plastic container • Form a fist around the EpiPen and pull off the blue safety release (cap) • Hold leg still and place orange end against the student's outer mid-thigh (with or without clothing) • Push down hard until a click is heard or felt and hold in place for 3 seconds • Remove EpiPen • Note the time the EpiPen is administered • Retain the used EpiPen to be handed to ambulance paramedics along with the time of administration. <p>OR</p> <p>Administer an Anapen® 500</p> <ul style="list-style-type: none"> • Pull off the black needle shield • Pull off grey safety cap (from the red button) • Place needle end firmly against the student's outer mid-thigh at 90 degrees (with or without clothing) • Press red button so it clicks and hold for 3 seconds • Remove Anapen® • Note the time the Anapen is administered • Retain the used Anapen to be handed to ambulance paramedics along with the time of administration. <p>OR</p> <p>Administer Jext 150 or 300</p>

	<ul style="list-style-type: none"> • Form fist around Jext and pull off yellow cap • Place black injector tip against outer-mid thigh (with or without clothing) • Push black tip firmly until a click is heard and hold in place for 3 seconds. • Remove Jext • Note the time the Jext device is administered. • The used adrenaline device must be handed to the ambulance paramedics along with the time of administration <p>OR</p> <p>Administer Neffy® 1mg or 2mg</p> <ul style="list-style-type: none"> • Hold the nasal spray with your thumb on the bottom of the plunger and a finger on either side of the nozzle. • Do not pull or push on the plunger. Do not test or prime (pre-spray). Each Neffy nasal spray contains only one spray. • Place the nozzle of the nasal spray into a nostril until fingers touch the nose. • For smaller nostrils, aim for the fingers to touch the nose. • Keep the nozzle pointed towards the forehead. Do not angle the nozzle of the nasal spray to the inner or outer walls of the nose. • Press the plunger up firmly until the dose is administered and it sprays into the nostril. • Note the time the Neffy device is administered. • The used adrenaline device must be handed to the ambulance paramedics along with the time of administration
3.	Call an ambulance (000)
4.	If there is no improvement or severe symptoms progress (as described in the ASCIA Action Plan for Anaphylaxis (RED)), further adrenaline doses may be administered every 5 minutes, if other adrenaline autoinjectors are available.
5.	Contact the student's emergency contacts.
6.	The principal or a staff member allocated to do so must contact the Incident Support and Operations Centre (ISOC) on 1800 126 126 to report 'High' or Extreme' severity incidents to report the incident. Incidents assessed as 'Low' or 'Medium' can be reported directly into EduSafe Plus by the principal or their allocated staff member.

If a student appears to be having a severe allergic reaction but has not been previously diagnosed with an allergy or being at risk of anaphylaxis, school staff should follow steps 2 – 5 as above.

For first time anaphylactic reactions, the school's general use adrenaline autoinjector device must be used. If the general use device is not immediately available in an anaphylaxis emergency, staff may use another student's adrenaline device, including the Epipen®, Anapen®, Jext® or Neffy® device. This may save a life. If another student's adrenaline device is used in an anaphylaxis emergency, the school must notify the parents of the student whose device was used and immediately replace the device.

Where possible, schools should consider using the correctly dosed adrenaline device depending on the weight of the student. However, in an emergency if there is no other option available, any device should be administered to the student.

Communication Plan

This policy will be publicly available on Carrum Primary School's website so that parents and other members of the school community can easily access information about Carrum Primary School's anaphylaxis management procedures. The parents and carers of students who are enrolled at Carrum

Primary School and are identified as being at risk of anaphylaxis will also be provided with a copy of this policy.

The principal is responsible for ensuring that all relevant staff, including casual relief staff, canteen staff and volunteers are aware of this policy and Carrum Primary School's procedures for anaphylaxis management. We will communicate this policy through staff induction and training materials, and twice-yearly staff briefings. Casual relief staff and volunteers who are responsible for the care and/or supervision of students who are identified as being at risk of anaphylaxis will also receive a verbal briefing on this policy (on annual induction), their role in responding to an anaphylactic reaction and where required, the identity of students at risk.

We will also communicate this policy through:

- Usual school community communication platform (Compass)
- Staff manual
- School Council induction materials
- Annual communication to school community
- Parent information night
- Website

The Principal is also responsible for ensuring relevant staff are trained and briefed in anaphylaxis management, consistent with the department's [Anaphylaxis Guidelines](#).

Staff training

The principal will ensure that the following school staff are appropriately trained in anaphylaxis management:

- School staff who conduct classes attended by students who are at risk of anaphylaxis
- School staff who conduct specialist classes, admin staff, first aiders and any other member of school staff as required by the principal based on a risk assessment.

Staff who are required to undertake training must have completed:

- an approved face-to-face anaphylaxis management training course in the last three years, or
- an approved online anaphylaxis management training course in the last two years.

Carrum Primary School uses the following training courses ASCIA Anaphylaxis e-Training course for Victorian Schools and/or Course in First Aid Management of Anaphylaxis 22578VIC formally 22300VIC.

Staff are also required to attend a briefing on anaphylaxis management and this policy at least twice per year (with the first briefing to be held at the beginning of the school year), facilitated by a staff member who has successfully completed an anaphylaxis management course within the last 2 years including the principal or School Anaphylaxis Supervisor. Each briefing should address:

- the school's anaphylaxis management policy
- the causes, symptoms and treatment of anaphylaxis
- the identities of students diagnosed as being at risk of anaphylaxis, their allergens and the location of their Individual Anaphylaxis Management Plans and their medication/s
- discussion on staff anaphylaxis training and renewal
- how to use an adrenaline device, including hands-on practice with an adrenaline device trainer device (which does not contain adrenaline)
- the school's general first aid and emergency procedures
- the location of adrenaline autoinjector devices prescribed for individual students that have been purchased by their family
- the location of adrenaline devices that the school has purchased for general use

- how to access on-going support and training.

When a new student enrolls at Carrum Primary School who is at risk of anaphylaxis, the principal will develop an interim plan in consultation with the student’s parents and ensure that appropriate staff are trained and briefed as soon as possible.

A record of staff training courses and briefings will be maintained in our school administration Occupational Health and Safety folder under First Aid and Infection Control – Staff training and on our online Emergency Management Plan.

The principal will ensure that while students at risk of anaphylaxis are under the care or supervision of the school outside of normal class activities, including in the school yard, at camps and excursions, or at special event days, there is a sufficient number of school staff present who have been trained in anaphylaxis management.

FURTHER INFORMATION AND RESOURCES

- The department’s Policy and Advisory Library (PAL):
 - [Anaphylaxis](#)
 - Allergies
 - First Aid for Students and Staff
 - Health Care Needs
 - Managing Reporting School Incidents (Including Emergencies)
 - Medication
- [Allergy & Anaphylaxis Australia](#)
- ASCIA Guidelines: [Schooling and childcare](#)
- [Hero HQ Anaphylaxis Management Training](#)
- https://allergyfacts.org.au/__interest/anaphylaxis/
- Royal Children’s Hospital: [Allergy and immunology](#)
- The school’s local policies: Health Care Needs, First Aid Policy, Medication

POLICY REVIEW AND APPROVAL

Policy last reviewed	12/March/2026
Approved by	Principal
Next scheduled review date	March 2027

The principal and delegate will complete the department’s Annual Risk Management Checklist for anaphylaxis management to assist with the evaluation and review of this policy and the support provided to students at risk of anaphylaxis.